

# Application rates for C-BIO CPS

Crop	Foliar dose (L/Ha)	Fertigation (L/Ha)	Application
Olive and Vine	1-1.5	1.5-2	2-4 application
Table grapes	1-2	2-3	10-15 days
Strawberry, raspberry and blueberries	1-2	2-3	7-10 days
Root and Bulbous Vegetables	0.75-1.5	1.5-3	10-15
Vegetables in the nursery	0.75-1.5	1.5-2	Apply in seedbed at a general dose of 100-150ml/hl
Ornamental Plants	1-1.5	1.5-2	3 intervals every 15 days from the vegetative stage
Grassland (Pasture/silage)	3-5	5-7	Apply between February – September
Turf	5-20	20-25	2-4 weeks
Potatoes	0.75- 1.5	1.5-2.5	7-14 days
Sugar Beet	2-5	4-6	Growth Stage 12-15, Growth Stage 19, Growth stage 31-33
Wheat and Barley	1-2	1.5-3	Growth stage 12-15, T0, T1
Oilseed Rape	2-3	3-5	Growth stage 1,3 and 5
Green house vegetables	1-2	2-3	7-10 days, 2-5 application
Open-Air Vegetables	0.75-1.5	1.5-2	10-15 days, 2-3
Leafy vegetables	1-3	2.5-3.5	10-15 days
Citrus Fruit	1-2	2-3	10-15 days
Stone and shell fruit	1-1.5	2-2.5	10-15 days
Shake the container well before use. Dilute the required quantity of C-Bio in a minimum of 200 liters of water per Hectare			